

Works Cited

- Holte, J., & Talley, L. (2008). A New Algorithm for Finding Mixed Layer Depths with Applications to Argo Data and Subantarctic Mode Water Formation. *Journal of Atmospheric and Oceanic Technology*. Retrieved from https://journals.ametsoc.org/view/journals/atot/26/9/2009jtecho543_1.xml?tab_body=fulltext-display
- Horrocks, L. A., O'Carroll, A., Candy, B., Nightingale, T. J., & Harris, A. R. (2003). Parameterizations of the ocean skin effect and implications for satellite-based measurement of sea-surface temperature. *Journal of Geophysical Research*. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/full/10.1029/2002JC001503>
- Levitus, S. (1982). *Climatological Atlas of the World Ocean*. Rockville, Md. Retrieved from https://www.google.com/books/edition/Climatological_Atlas_of_the_World_Ocean/_x0IAQAAIAAJ?hl=en&gbpv=0
- Wick, G., & Castro, S. (2020). Assessment of Extreme Diurnal Warming in Operational Geosynchronous Satellite Sea Surface Temperature Products. *Remote Sensing*, 12(22), 3771. Retrieved from <https://www.mdpi.com/2072-4292/12/22/3771>
- Wurl, O., Bird, K., Cunliffe, M., Landing, W., Miller, U., Mustaffa, N., . . . Zappa, C. (2018). Warming and Inhibition of Salinization at the Ocean's Surface by Cyanobacteria. *Geophysical Research Letters*, 45(9), 4230-4237. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/full/10.1029/2018GL077946>